

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/54, 9/12, C12P 21/00, A61K 38/45, C12Q 1/68, C07K 16/40</b>		<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 99/11794</b>
			<b>(43) International Publication Date:</b> 11 March 1999 (11.03.99)
<b>(21) International Application Number:</b> PCT/US98/18524 <b>(22) International Filing Date:</b> 4 September 1998 (04.09.98) <b>(30) Priority Data:</b> 08/923,469 4 September 1997 (04.09.97) US <b>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application</b> US 08/923,469 (CIP) Filed on 4 September 1997 (04.09.97) <b>(71) Applicant (for all designated States except US):</b> INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> YUE, Henry [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). LAL, Preeti [IN/US]; 2382 Lass Drive, Santa Clara, CA 95054 (US). CORLEY, Neil, C. [US/US]; 1240 Dale Avenue #30, Mountain View, CA 94040 (US). <b>(74) Agents:</b> BILLINGS, Lucy, J. et al.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US).			<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> NEW VRK1 KINASE			
<b>(57) Abstract</b> <p>The invention provides a new VRK1 kinase (NVRK1) and polynucleotides which identify and encode NVRK1. The invention also provides expression vectors, host cells, agonists, antibodies and antagonists. The invention also provides methods for treating disorders associated with expression of NVRK1.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## New VRK1 Kinase

### TECHNICAL FIELD

5 This invention relates to nucleic acid and amino acid sequences of a new VRK1 kinase and to the use of these sequences in the diagnosis, prevention, and treatment of diseases associated with cell proliferation.

### BACKGROUND OF THE INVENTION

10 Kinases regulate many different cell proliferation, differentiation, and signaling processes by adding phosphate groups to proteins. Uncontrolled signaling has been implicated in a variety of disease conditions including inflammation, cancer, arteriosclerosis, and psoriasis. Reversible protein phosphorylation is the main strategy for controlling activities of eukaryotic cells. It is estimated that more than 1000 of the 10,000 proteins active  
15 in a typical mammalian cell are phosphorylated. The high energy phosphate which drives activation is generally transferred from adenosine triphosphate molecules (ATP) to a particular protein by protein kinases and removed from that protein by protein phosphatases. Phosphorylation occurs in response to extracellular signals (hormones, neurotransmitters, growth and differentiation factors, etc), cell cycle checkpoints, and environmental or  
20 nutritional stresses and is roughly analogous to turning on a molecular switch. When the switch goes on, the appropriate protein kinase activates a metabolic enzyme, regulatory protein, receptor, cytoskeletal protein, ion channel or pump, or transcription factor.

Protein tyrosine kinases, PTKs, specifically phosphorylate tyrosine residues on their target proteins and may be divided into transmembrane, receptor PTKs and  
25 nontransmembrane, non-receptor PTKs. Transmembrane protein-tyrosine kinases are receptors for most growth factors. Binding of growth factor to the receptor activates the transfer of a phosphate group from ATP to selected tyrosine side chains of the receptor and other specific proteins. Growth factors (GF) associated with receptor PTKs include; epidermal GF, platelet-derived GF, fibroblast GF, hepatocyte GF, insulin and insulin-like  
30 GFs, nerve GF, vascular endothelial GF, and macrophage colony stimulating factor.

Non-receptor PTKs lack transmembrane regions and, instead, form complexes with the intracellular regions of cell surface receptors. Some of the receptors that function through non-receptor PTKs include those for cytokines and hormones (growth hormone and prolactin)

and antigen-specific receptors on the surface of T and B lymphocytes. The protein products of oncogenes and many growth-factor receptors have protein kinase activities that phosphorylate tyrosine.

An additional family of proteins kinases phosphorylates serine and threonine and are found to regulate hormone activity by increasing the cellular concentration of second messengers such as cyclic AMP, diacylglycerol, or  $\text{Ca}^{2+}$ . In turn, these second messengers activate protein kinases that phosphorylate serine and threonine residues of various proteins. An example is the vaccinia virus, B1R kinase: a 36-kDa protein induced in early stages of infection. Phosphorylation of the vaccinia virus B1R yeilds multiple forms with isoelectric points (pI) of 5.5, 5.7, 5.9, and 6.3. The tryptic peptides derived from these isoforms produce almost identical high-pressure liquid chromatography profiles. The similarity of these profiles strongly suggest that the 36-kDa protein was modified by phosphorylation of at least four sites, which have been identified as threonine residues. (Beaud G. et al. (1995) J. Virol. 69(3):1819-1826).

The discovery of a new VRK1 kinase and the polynucleotides encoding it satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention and treatment of diseases associated with cell proliferation, in particular, immune responses and cancers.

## SUMMARY OF THE INVENTION

The invention features a substantially purified polypeptide, new VRK1 kinase, having the amino acid sequence shown in SEQ ID NO:1, or fragments thereof.

The invention further provides an isolated and substantially purified polynucleotide sequence encoding the polypeptide comprising the amino acid sequence of SEQ ID NO:1 or fragments thereof and a composition comprising said polynucleotide sequence. The invention also provides a polynucleotide sequence which hybridizes under stringent conditions to the polynucleotide sequence encoding the amino acid sequence SEQ ID NO:1, or fragments of said polynucleotide sequence. The invention further provides a polynucleotide sequence comprising the complement of the polynucleotide sequence encoding the amino acid sequence of SEQ ID NO:1, or fragments or variants of said polynucleotide sequence.

The invention also provides an isolated and purified sequence comprising SEQ ID NO:2 or variants thereof. In addition, the invention provides a polynucleotide sequence which hybridizes under stringent conditions to the polynucleotide sequence of SEQ ID NO:2.

The invention also provides a polynucleotide sequence comprising the complement of SEQ ID NO:2, or fragments or variants thereof.

The present invention further provides an expression vector containing at least a fragment of any of the claimed polynucleotide sequences. In yet another aspect, the expression vector containing the polynucleotide sequence is contained within a host cell.

The invention also provides a method for producing a polypeptide comprising the amino acid sequence of SEQ ID NO:1 or a fragment thereof, the method comprising the steps of: a) culturing the host cell containing an expression vector containing at least a fragment of the polynucleotide sequence encoding NVRK1 under conditions suitable for the expression of the polypeptide; and b) recovering the polypeptide from the host cell culture.

The invention also provides a pharmaceutical composition comprising a substantially purified NVRK1 having the amino acid sequence of SEQ ID NO:1 in conjunction with a suitable pharmaceutical carrier.

The invention also provides a purified antagonist of the polypeptide of SEQ ID NO:1. In one aspect the invention provides a purified antibody which binds to a polypeptide comprising the amino acid sequence of SEQ ID NO:1.

Still further, the invention provides a purified agonist of the polypeptide of SEQ ID NO:1.

The invention also provides a method for treating or preventing an immune response comprising administering to a subject in need of such treatment an effective amount of an antagonist to NVRK1.

The invention also provides a method for treating a cancer comprising administering to a subject in need of such treatment an effective amount of an antagonist to NVRK1.

The invention also provides a method for detecting a polynucleotide which encodes NVRK1 in a biological sample comprising the steps of: a) hybridizing the complement of the polynucleotide sequence which encodes SEQ ID NO:1 to nucleic acid material of a biological sample, thereby forming a hybridization complex; and b) detecting the hybridization complex, wherein the presence of the complex correlates with the presence of a polynucleotide encoding NVRK1 in the biological sample. In one aspect the nucleic acid material of the biological sample is amplified by the polymerase chain reaction prior to hybridization.

## BRIEF DESCRIPTION OF THE FIGURES

Figures 1A, 1B, 1C, 1D, and 1E shows the amino acid sequence (SEQ ID NO:1) and nucleic acid sequence (SEQ ID NO:2) of NVRK1. The alignment was produced using MacDNASIS PRO™ software (Hitachi Software Engineering Co. Ltd. San Bruno, CA).

Figures 2A and 2B show the amino acid sequence alignments among NVRK1 (SEQ ID NO:1), and vaccinia virus B1R kinase (GI 222701; SEQ ID NO:3), produced using the multisequence alignment program of DNASTAR™ software (DNASTAR Inc, Madison WI).

Figures 3A and 3B show the hydrophobicity plots for NVRK1 (SEQ ID NO: 1) and vaccinia virus B1R kinase (SEQ ID NO:3), respectively ; the positive X axis reflects amino acid position, and the negative Y axis, hydrophobicity (MacDNASIS PRO software).

Figure 4 shows an isoelectric plot for NVRK1 (SEQ ID NO: 1) produced using MacDNASIS PRO software).

## DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular methodology, protocols, cell lines, vectors, and reagents described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms “a”, “an”, and “the” include plural reference unless the context clearly dictates otherwise. Thus, for example, reference to “a host cell” includes a plurality of such host cells, reference to the “antibody” is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods, devices, and materials are now described. All publications mentioned herein are incorporated herein by reference for the purpose of describing and disclosing the cell lines, vectors, and methodologies which are reported in the publications which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

**DEFINITIONS**

NVRK1, as used herein, refers to the amino acid sequences of substantially purified NVRK1 obtained from any species, particularly mammalian, including bovine, ovine, porcine, murine, equine, and preferably human, from any source whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist", as used herein, refers to a molecule which, when bound to NVRK1, increases or prolongs the duration of the effect of NVRK1. Agonists may include proteins, nucleic acids, carbohydrates, or any other molecules which bind to and modulate the effect of NVRK1.

An "allele" or "allelic sequence", as used herein, is an alternative form of the gene encoding NVRK1. Alleles may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or polypeptides whose structure or function may or may not be altered. Any given natural or recombinant gene may have none, one, or many allelic forms. Common mutational changes which give rise to alleles are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

"Altered" nucleic acid sequences encoding NVRK1 as used herein include those with deletions, insertions, or substitutions of different nucleotides resulting in a polynucleotide that encodes the same or a functionally equivalent NVRK1. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding NVRK1, and improper or unexpected hybridization to alleles, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding NVRK1. The encoded protein may also be "altered" and contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent NVRK1. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues as long as the biological or immunological activity of NVRK1 is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid; positively charged amino acids may include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values may include leucine, isoleucine, and valine, glycine and alanine, asparagine and glutamine, serine and threonine, and phenylalanine and tyrosine.

“Amino acid sequence” as used herein refers to an oligopeptide, peptide, polypeptide, or protein sequence, and fragment thereof, and to naturally occurring or synthetic molecules. Fragments of NVRK1 are preferably about 5 to about 15 amino acids in length and retain the biological activity or the immunological activity of NVRK1. Where “amino acid sequence” is recited herein to refer to an amino acid sequence of a naturally occurring protein molecule, amino acid sequence, and like terms, are not meant to limit the amino acid sequence to the complete, native amino acid sequence associated with the recited protein molecule.

“Amplification” as used herein refers to the production of additional copies of a nucleic acid sequence and is generally carried out using polymerase chain reaction (PCR) technologies well known in the art (Dieffenbach, C.W. and G.S. Dveksler (1995) PCR Primer, a Laboratory Manual, Cold Spring Harbor Press, Plainview, NY).

The term “antagonist” as used herein, refers to a molecule which, when bound to NVRK1, decreases the amount or the duration of the effect of the biological or immunological activity of NVRK1. Antagonists may include proteins, nucleic acids, carbohydrates, antibodies or any other molecules which decrease the effect of NVRK1.

As used herein, the term “antibody” refers to intact molecules as well as fragments thereof, such as Fa, F(ab')<sub>2</sub>, and Fv, which are capable of binding the epitopic determinant. Antibodies that bind NVRK1 polypeptides can be prepared using intact polypeptides or fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal can be derived from the translation of RNA or synthesized chemically and can be conjugated to a carrier protein, if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin and thyroglobulin, keyhole limpet hemocyanin. The coupled peptide is then used to immunize the animal (e.g., a mouse, a rat, or a rabbit).

The term “antigenic determinant”, as used herein, refers to that fragment of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to a given region or three-dimensional structure on the protein; these regions or structures are referred to as antigenic determinants. An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term “antisense”, as used herein, refers to any composition containing nucleotide sequences which are complementary to a specific DNA or RNA sequence. The term



"antisense strand" is used in reference to a nucleic acid strand that is complementary to the "sense" strand. Antisense molecules include peptide nucleic acids and may be produced by any method including synthesis or transcription. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form duplexes and block either transcription or translation. The designation "negative" is sometimes used in reference to the antisense strand, and "positive" is sometimes used in reference to the sense strand.

The term "biologically active", as used herein, refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" refers to the capability of the natural, recombinant, or synthetic NVRK1, or any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The terms "complementary" or "complementarity", as used herein, refer to the natural binding of polynucleotides under permissive salt and temperature conditions by base-pairing. For example, the sequence "A-G-T" binds to the complementary sequence "T-C-A". Complementarity between two single-stranded molecules may be "partial", in which only some of the nucleic acids bind, or it may be complete when total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of hybridization between nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acids strands and in the design and use of PNA molecules.

A "composition comprising a given polynucleotide sequence" as used herein refers broadly to any composition containing the given polynucleotide sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding NVRK1 (SEQ ID NO:1) or fragments thereof (e.g., SEQ ID NO:2 and fragments thereof) may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., SDS) and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus", as used herein, refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, has been extended using XL-PCR™ (Perkin Elmer, Norwalk, CT) in the 5' and/or the 3' direction and resequenced, or has been assembled from

the overlapping sequences of more than one Incyte Clone using a computer program for fragment assembly (e.g., GELVIEW™ Fragment Assembly system, GCG, Madison, WI). Some sequences have been both extended and assembled to produce the consensus sequence.

The term “correlates with expression of a polynucleotide”, as used herein, indicates that the detection of the presence of ribonucleic acid that is similar to SEQ ID NO:2 by northern analysis is indicative of the presence of mRNA encoding NVRK1 in a sample and thereby correlates with expression of the transcript from the polynucleotide encoding the protein.

A “deletion”, as used herein, refers to a change in the amino acid or nucleotide sequence and results in the absence of one or more amino acid residues or nucleotides.

The term “derivative”, as used herein, refers to the chemical modification of a nucleic acid encoding or complementary to NVRK1 or the encoded NVRK1. Such modifications include, for example, replacement of hydrogen by an alkyl, acyl, or amino group. A nucleic acid derivative encodes a polypeptide which retains the biological or immunological function of the natural molecule. A derivative polypeptide is one which is modified by glycosylation, pegylation, or any similar process which retains the biological or immunological function of the polypeptide from which it was derived.

The term “homology”, as used herein, refers to a degree of complementarity. There may be partial homology or complete homology (i.e., identity). A partially complementary sequence that at least partially inhibits an identical sequence from hybridizing to a target nucleic acid is referred to using the functional term “substantially homologous.” The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or northern blot, solution hybridization and the like) under conditions of low stringency. A substantially homologous sequence or hybridization probe will compete for and inhibit the binding of a completely homologous sequence to the target sequence under conditions of low stringency. This is not to say that conditions of low stringency are such that non-specific binding is permitted; low stringency conditions require that the binding of two sequences to one another be a specific (i.e., selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% identity). In the absence of non-specific binding, the probe will not hybridize to the second non-complementary target sequence.

Human artificial chromosomes (HACs) are linear microchromosomes which may

contain DNA sequences of 10K to 10M in size and contain all of the elements required for stable mitotic chromosome segregation and maintenance (Harrington, J.J. et al. (1997) Nat Genet. 15:345-355). The term "humanized antibody", as used herein, refers to antibody molecules in which amino acids have been replaced in the non-antigen binding regions in order to more closely resemble a human antibody, while still retaining the original binding ability.

The term "hybridization", as used herein, refers to any process by which a strand of nucleic acid binds with a complementary strand through base pairing.

The term "hybridization complex", as used herein, refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary G and C bases and between complementary A and T bases; these hydrogen bonds may be further stabilized by base stacking interactions. The two complementary nucleic acid sequences hydrogen bond in an antiparallel configuration. A hybridization complex may be formed in solution (e.g., C<sub>0</sub>t or R<sub>0</sub>t analysis) or between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

An "insertion" or "addition", as used herein, refers to a change in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively, as compared to the naturally occurring molecule.

"Microarray" refers to an array of distinct polynucleotides or oligonucleotides synthesized on a substrate, such as paper, nylon or other type of membrane, filter, chip, glass slide, or any other suitable solid support.

The term "modulate", as used herein, refers to a change in the activity of NVRK1. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional or immunological properties of NVRK1.

"Nucleic acid sequence" as used herein refers to an oligonucleotide, nucleotide, or polynucleotide, and fragments thereof, and to DNA or RNA of genomic or synthetic origin which may be single- or double-stranded, and represent the sense or antisense strand.

"Fragments" are those nucleic acid sequences which are greater than 60 nucleotides in length, and most preferably includes fragments that are at least 100 nucleotides or at least 1000 nucleotides, and at least 10,000 nucleotides in length.

The term "oligonucleotide" refers to a nucleic acid sequence of at least about 6

nucleotides to about 60 nucleotides, preferably about 15 to 30 nucleotides, and more preferably about 20 to 25 nucleotides, which can be used in PCR amplification or a hybridization assay, or a microarray. As used herein, oligonucleotide is substantially equivalent to the terms "amplimers", "primers", "oligomers", and "probes", as commonly defined in the art.

"Peptide nucleic acid", PNA as used herein, refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least five nucleotides in length linked to a peptide backbone of amino acid residues which ends in lysine. The terminal lysine confers solubility to the composition. PNAs may be pegylated to extend their lifespan in the cell where they preferentially bind complementary single stranded DNA and RNA and stop transcript elongation (Nielsen, P.E. et al. (1993) Anticancer Drug Des. 8:53-63).

The term "portion", as used herein, with regard to a protein (as in "a portion of a given protein") refers to fragments of that protein. The fragments may range in size from five amino acid residues to the entire amino acid sequence minus one amino acid. Thus, a protein "comprising at least a portion of the amino acid sequence of SEQ ID NO:1" encompasses the full-length NVRK1 and fragments thereof.

The term "sample", as used herein, is used in its broadest sense. A biological sample suspected of containing nucleic acid encoding NVRK1, or fragments thereof, or NVRK1 itself may comprise a bodily fluid, extract from a cell, chromosome, organelle, or membrane isolated from a cell, a cell, genomic DNA, RNA, or cDNA (in solution or bound to a solid support, a tissue, a tissue print, and the like).

The terms "specific binding" or "specifically binding", as used herein, refers to that interaction between a protein or peptide and an agonist, an antibody and an antagonist. The interaction is dependent upon the presence of a particular structure (i.e., the antigenic determinant or epitope) of the protein recognized by the binding molecule. For example, if an antibody is specific for epitope "A", the presence of a protein containing epitope A (or free, unlabeled A) in a reaction containing labeled "A" and the antibody will reduce the amount of labeled A bound to the antibody.

The terms "stringent conditions" or "stringency", as used herein, refer to the conditions for hybridization as defined by the nucleic acid, salt, and temperature. These conditions are well known in the art and may be altered in order to identify or detect identical or related polynucleotide sequences. Numerous equivalent conditions comprising either low or high stringency depend on factors such as the length and nature of the sequence (DNA, RNA, base

composition), nature of the target (DNA, RNA, base composition), milieu (in solution or immobilized on a solid substrate), concentration of salts and other components (e.g., formamide, dextran sulfate and/or polyethylene glycol), and temperature of the reactions (within a range from about 5°C below the melting temperature of the probe to about 20°C to 25°C below the melting temperature). One or more factors may be varied to generate conditions of either low or high stringency different from, but equivalent to, the above listed conditions.

The term “substantially purified”, as used herein, refers to nucleic or amino acid sequences that are removed from their natural environment, isolated or separated, and are at least 60% free, preferably 75% free, and most preferably 90% free from other components with which they are naturally associated.

A “substitution”, as used herein, refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

“Transformation”, as defined herein, describes a process by which exogenous DNA enters and changes a recipient cell. It may occur under natural or artificial conditions using various methods well known in the art. Transformation may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method is selected based on the type of host cell being transformed and may include, but is not limited to, viral infection, electroporation, heat shock, lipofection, and particle bombardment. Such “transformed” cells include stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome. They also include cells which transiently express the inserted DNA or RNA for limited periods of time.

A “variant” of NVRK1, as used herein, refers to an amino acid sequence that is altered by one or more amino acids. The variant may have “conservative” changes, wherein a substituted amino acid has similar structural or chemical properties, e.g., replacement of leucine with isoleucine. More rarely, a variant may have “nonconservative” changes, e.g., replacement of a glycine with a tryptophan. Analogous minor variations may also include amino acid deletions or insertions, or both. Guidance in determining which amino acid residues may be substituted, inserted, or deleted without abolishing biological or immunological activity may be found using computer programs well known in the art, for example, DNASTAR software.

**THE INVENTION**

The invention is based on the discovery of a new VRK1 kinase (hereinafter referred to as "NVRK1"), the polynucleotides encoding NVRK1, and the use of these compositions for the diagnosis, prevention, or treatment of cancer and immune respons.

5 Nucleic acids encoding the NVRK1 of the present invention were first identified in Incyte Clone 1543990 from the prostate cDNA library (PROSTUT04) using a computer search for amino acid sequence alignments. A consensus sequence, SEQ ID NO:2, was derived from the following overlapping and/or extended nucleic acid sequences: Incyte Clones 1543990 (PROSTUT04), 1694220, 1696636 (COLNNOT23), and 2221838  
10 (LUNGNOT18).

In one embodiment, the invention encompasses a polypeptide comprising the amino acid sequence of SEQ ID NO:1, as shown in Figs. 1A, 1B, 1C, 1D, and 1E. NVRK1 is 396 amino acids in length and has one potential cAMP- and cGMP-dependent protein kinase phosphorylation site at K<sub>147</sub>; four potential casein kinase II phosphorylation sites at S<sub>284</sub>, S<sub>333</sub>,  
15 S<sub>342</sub>, and T<sub>378</sub>; ten potential protein kinase C phosphorylation sites at residues S<sub>14</sub>, T<sub>102</sub>, S<sub>125</sub>, S<sub>150</sub>, S<sub>158</sub>, S<sub>239</sub>, T<sub>305</sub>, T<sub>312</sub>, T<sub>355</sub>, and T<sub>390</sub>; four potential tyrosine phosphorylation sites at residues R<sub>160</sub>, K<sub>188</sub>, R<sub>241</sub>, and K<sub>295</sub>; and one protein kinase signature and profile site at I<sub>43</sub>. As shown in Figure 2, NVRK1 has chemical and structural homology with vaccinia virus B1R kinase (GI 222701; SEQ ID NO:3). In particular, NVRK1 and vaccinia virus B1R kinase  
20 share 43% identity and the signature and profile site at I<sub>43</sub>, the tyrosine phosphorylation site at R<sub>241</sub>, and phosphorylation sites at K<sub>188</sub>, S<sub>239</sub>, and R<sub>241</sub>. As illustrated by Figs. 3A and 3B, NVRK1 and vaccinia virus B1R kinase have rather similar hydrophobicity plots and isoelectric points (Figure 4). Northern analysis shows the expression of NVRK1 in various cDNA libraries, at least 47% of which are immortalized or cancerous, at least 38% of which  
25 involve immune response, and at least 22% of which involve fetal/infant development.

The invention also encompasses NVRK1 variants. A preferred NVRK1 variant is one having at least 80%, and more preferably at least 90%, amino acid sequence identity to the NVRK1 amino acid sequence (SEQ ID NO:1) and which retains at least one biological, immunological or other functional characteristic or activity of NVRK1. A most preferred  
30 NVRK1 variant is one having at least 95% amino acid sequence identity to SEQ ID NO:1.

The invention also encompasses polynucleotides which encode NVRK1. Accordingly, any nucleic acid sequence which encodes the amino acid sequence of NVRK1 can be used to produce recombinant molecules which express NVRK1. In a particular

embodiment, the invention encompasses the polynucleotide comprising the nucleic acid sequence of SEQ ID NO:2 as shown in Figure 1.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of nucleotide sequences encoding NVRK1, some bearing minimal  
5 homology to the nucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of nucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the nucleotide sequence of naturally occurring NVRK1, and all such variations are to be  
10 considered as being specifically disclosed.

Although nucleotide sequences which encode NVRK1 and its variants are preferably capable of hybridizing to the nucleotide sequence of the naturally occurring NVRK1 under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding NVRK1 or its derivatives possessing a substantially different codon  
15 usage. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding NVRK1 and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties,  
20 such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences, or fragments thereof, which encode NVRK1 and its derivatives, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents that are well known in the art. Moreover, synthetic chemistry may  
25 be used to introduce mutations into a sequence encoding NVRK1 or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed nucleotide sequences, and in particular, those shown in SEQ ID NO:2, under various conditions of stringency as taught in Wahl, G.M. and S.L. Berger (1987; Methods Enzymol. 152:399-407) and Kimmel, A.R. (1987; Methods Enzymol. 152:507-  
30 511).

Methods for DNA sequencing which are well known and generally available in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, Sequenase® (US

Biochemical Corp, Cleveland, OH), Taq polymerase (Perkin Elmer), thermostable T7 polymerase (Amersham, Chicago, IL), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE Amplification System marketed by Gibco/BRL (Gaithersburg, MD). Preferably, the process is automated with machines such as the Hamilton Micro Lab 2200 (Hamilton, Reno, NV), Peltier Thermal Cycler (PTC200; MJ Research, Watertown, MA) and the ABI Catalyst and 373 and 377 DNA Sequencers (Perkin Elmer).

The nucleic acid sequences encoding NVRK1 may be extended utilizing a partial nucleotide sequence and employing various methods known in the art to detect upstream sequences such as promoters and regulatory elements. For example, one method which may be employed, "restriction-site" PCR, uses universal primers to retrieve unknown sequence adjacent to a known locus (Sarkar, G. (1993) PCR Methods Applic. 2:318-322). In particular, genomic DNA is first amplified in the presence of primer to a linker sequence and a primer specific to the known region. The amplified sequences are then subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

Inverse PCR may also be used to amplify or extend sequences using divergent primers based on a known region (Triglia, T. et al. (1988) Nucleic Acids Res. 16:8186). The primers may be designed using commercially available software such as OLIGO 4.06 Primer Analysis software (National Biosciences Inc., Plymouth, MN), or another appropriate program, to be 22-30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68°-72° C. The method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

Another method which may be used is capture PCR which involves PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial chromosome DNA (Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119). In this method, multiple restriction enzyme digestions and ligations may also be used to place an engineered double-stranded sequence into an unknown fragment of the DNA molecule before performing PCR.

Another method which may be used to retrieve unknown sequences is that of Parker, J.D. et al. (1991; Nucleic Acids Res. 19:3055-3060). Additionally, one may use PCR, nested



primers, and PromoterFinder™ libraries to walk genomic DNA (Clontech, Palo Alto, CA). This process avoids the need to screen libraries and is useful in finding intron/exon junctions.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. Also, random-primed libraries are preferable, in that they will contain more sequences which contain the 5' regions of genes. Use of a randomly primed library may be especially preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different fluorescent dyes (one for each nucleotide) which are laser activated, and detection of the emitted wavelengths by a charge coupled device camera. Output/light intensity may be converted to electrical signal using appropriate software (e.g. Genotyper™ and Sequence Navigator™, Perkin Elmer) and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for the sequencing of small pieces of DNA which might be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode NVRK1 may be used in recombinant DNA molecules to direct expression of NVRK1, fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced, and these sequences may be used to clone and express NVRK1.

As will be understood by those of skill in the art, it may be advantageous to produce NVRK1-encoding nucleotide sequences possessing non-naturally occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce an RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter NVRK1 encoding sequences for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or

expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, site-directed mutagenesis may be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, introduce mutations, and so forth.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding NVRK1 may be ligated to a heterologous sequence to encode a fusion protein. For example, to screen peptide libraries for inhibitors of NVRK1 activity, it may be useful to encode a chimeric NVRK1 protein that can be recognized by a commercially available antibody. A fusion protein may also be engineered to contain a cleavage site located between the NVRK1 encoding sequence and the heterologous protein sequence, so that NVRK1 may be cleaved and purified away from the heterologous moiety.

In another embodiment, sequences encoding NVRK1 may be synthesized, in whole or in part, using chemical methods well known in the art (see Caruthers, M.H. et al. (1980) Nucl. Acids Res. Symp. Ser. 215-223, Horn, T. et al. (1980) Nucl. Acids Res. Symp. Ser. 225-232). Alternatively, the protein itself may be produced using chemical methods to synthesize the amino acid sequence of NVRK1, or a fragment thereof. For example, peptide synthesis can be performed using various solid-phase techniques (Roberge, J.Y. et al. (1995) Science 269:202-204) and automated synthesis may be achieved, for example, using the ABI 431A Peptide Synthesizer (Perkin Elmer).

The newly synthesized peptide may be substantially purified by preparative high performance liquid chromatography (e.g., Creighton, T. (1983) Proteins, Structures and Molecular Principles, WH Freeman and Co., New York, NY). The composition of the synthetic peptides may be confirmed by amino acid analysis or sequencing (e.g., the Edman degradation procedure; Creighton, supra). Additionally, the amino acid sequence of NVRK1, or any part thereof, may be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

In order to express a biologically active NVRK1, the nucleotide sequences encoding NVRK1 or functional equivalents, may be inserted into appropriate expression vector, i.e., a vector which contains the necessary elements for the transcription and translation of the inserted coding sequence.

Methods which are well known to those skilled in the art may be used to construct

expression vectors containing sequences encoding NVRK1 and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. Such techniques are described in Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview, NY, and Ausubel, F.M. et al. (1989) Current Protocols in Molecular Biology, John Wiley & Sons, New York, NY.

A variety of expression vector/host systems may be utilized to contain and express sequences encoding NVRK1. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with virus expression vectors (e.g., baculovirus); plant cell systems transformed with virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. The invention is not limited by the host cell employed.

The "control elements" or "regulatory sequences" are those non-translated regions of the vector--enhancers, promoters, 5' and 3' untranslated regions--which interact with host cellular proteins to carry out transcription and translation. Such elements may vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the Bluescript® phagemid (Stratagene, LaJolla, CA) or pSport1™ plasmid (Gibco BRL) and the like may be used. The baculovirus polyhedrin promoter may be used in insect cells. Promoters or enhancers derived from the genomes of plant cells (e.g., heat shock, RUBISCO; and storage protein genes) or from plant viruses (e.g., viral promoters or leader sequences) may be cloned into the vector. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are preferable. If it is necessary to generate a cell line that contains multiple copies of the sequence encoding NVRK1, vectors based on SV40 or EBV may be used with an appropriate selectable marker.

In bacterial systems, a number of expression vectors may be selected depending upon the use intended for NVRK1. For example, when large quantities of NVRK1 are needed for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified may be used. Such vectors include, but are not limited to, the

multifunctional E. coli cloning and expression vectors such as Bluescript® (Stratagene), in which the sequence encoding NVRK1 may be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of  $\beta$ -galactosidase so that a hybrid protein is produced; pIN vectors (Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509); and the like. pGEX vectors (Promega, Madison, WI) may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems may be designed to include heparin, thrombin, or factor XA protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

In the yeast, Saccharomyces cerevisiae, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH may be used. For reviews, see Ausubel et al. (supra) and Grant et al. (1987) Methods Enzymol. 153:516-544.

In cases where plant expression vectors are used, the expression of sequences encoding NVRK1 may be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV may be used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J. 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used (Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; and Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105). These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. Such techniques are described in a number of generally available reviews (see, for example, Hobbs, S. or Murry, L.E. in McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York, NY; pp. 191-196.

An insect system may also be used to express NVRK1. For example, in one such system, Autographa californica nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in Spodoptera frugiperda cells or in Trichoplusia larvae. The sequences encoding NVRK1 may be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of NVRK1 will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses may then be used to infect, for example, S. frugiperda

cells or Trichoplusia larvae in which NVRK1 may be expressed (Engelhard, E.K. et al. (1994) Proc. Nat. Acad. Sci. 91:3224-3227).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding NVRK1 may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain a viable virus which is capable of expressing NVRK1 in infected host cells (Logan, J. and Shenk, T. (1984) Proc. Natl. Acad. Sci. 81:3655-3659). In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained and expressed in a plasmid. HACs of 6 to 10M are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes.

Specific initiation signals may also be used to achieve more efficient translation of sequences encoding NVRK1. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding NVRK1, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including the ATG initiation codon should be provided. Furthermore, the initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used, such as those described in the literature (Scharf, D. et al. (1994) Results Probl. Cell Differ. 20:125-162).

In addition, a host cell strain may be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be used to facilitate correct insertion, folding and/or function. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38),

are available from the American Type Culture Collection (ATCC; Bethesda, MD) and may be chosen to ensure the correct modification and processing of the foreign protein.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express NVRK1 may be transformed using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be proliferated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler, M. et al. (1977) Cell 11:223-32) and adenine phosphoribosyltransferase (Lowy, I. et al. (1980) Cell 22:817-23) genes which can be employed in tk<sup>-</sup> or apt<sup>-</sup> cells, respectively. Also, antimetabolite, antibiotic or herbicide resistance can be used as the basis for selection; for example, dhfr which confers resistance to methotrexate (Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. 77:3567-70); npt, which confers resistance to the aminoglycosides neomycin and G-418 (Colbere-Garapin, F. et al (1981) J. Mol. Biol. 150:1-14) and als or pat, which confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively (Murry, supra). Additional selectable genes have been described, for example, trpB, which allows cells to utilize indole in place of tryptophan, or hisD, which allows cells to utilize histinol in place of histidine (Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. 85:8047-51). Recently, the use of visible markers has gained popularity with such markers as anthocyanins,  $\beta$  glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, being widely used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system (Rhodes, C.A. et al. (1995) Methods Mol. Biol. 55:121-131).

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, its presence and expression may need to be confirmed. For example, if the sequence encoding NVRK1 is inserted within a marker gene sequence, transformed cells containing sequences encoding NVRK1 can be identified by the absence of marker gene

function. Alternatively, a marker gene can be placed in tandem with a sequence encoding NVRK1 under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

Alternatively, host cells which contain the nucleic acid sequence encoding NVRK1 and express NVRK1 may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein.

The presence of polynucleotide sequences encoding NVRK1 can be detected by DNA-DNA or DNA-RNA hybridization or amplification using probes or fragments or fragments of polynucleotides encoding NVRK1. Nucleic acid amplification based assays involve the use of oligonucleotides or oligomers based on the sequences encoding NVRK1 to detect transformants containing DNA or RNA encoding NVRK1.

A variety of protocols for detecting and measuring the expression of NVRK1, using either polyclonal or monoclonal antibodies specific for the protein are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on NVRK1 is preferred, but a competitive binding assay may be employed. These and other assays are described, among other places, in Hampton, R. et al. (1990; Serological Methods, a Laboratory Manual, APS Press, St Paul, MN) and Maddox, D.E. et al. (1983; J. Exp. Med. 158:1211-1216).

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding NVRK1 include oligolabeling, nick translation, end-labeling or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding NVRK1, or any fragments thereof may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits (Pharmacia & Upjohn, (Kalamazoo, MI); Promega (Madison WI); and U.S. Biochemical

Corp., Cleveland, OH). Suitable reporter molecules or labels, which may be used for ease of detection, include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding NVRK1 may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode NVRK1 may be designed to contain signal sequences which direct secretion of NVRK1 through a prokaryotic or eukaryotic cell membrane. Other constructions may be used to join sequences encoding NVRK1 to nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, WA). The inclusion of cleavable linker sequences such as those specific for Factor XA or enterokinase (Invitrogen, San Diego, CA) between the purification domain and NVRK1 may be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing NVRK1 and a nucleic acid encoding 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification on IMAC (immobilized metal ion affinity chromatography as described in Porath, J. et al. (1992, Prot. Exp. Purif. 3: 263-281) while the enterokinase cleavage site provides a means for purifying NVRK1 from the fusion protein. A discussion of vectors which contain fusion proteins is provided in Kroll, D.J. et al. (1993; DNA Cell Biol. 12:441-453).

In addition to recombinant production, fragments of NVRK1 may be produced by direct peptide synthesis using solid-phase techniques Merrifield J. (1963) J. Am. Chem. Soc. 85:2149-2154). Protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer). Various fragments of NVRK1 may be chemically synthesized separately and combined using chemical methods to produce the full length molecule.



## THERAPEUTICS

Chemical and structural homology exists between NVRK1 and the vaccinia virus B1R kinase (GI 1827452) which is associated with viral diseases and carcinogenesis. Therefore, NVRK1 appears to play a role in diseases associated with cell proliferation. Northern analysis shows that the expression of NVRK1 is associated with cancer and immune response.

Therefore, in one embodiment, an antagonist of NVRK1 may be administered to a subject to prevent or treat an immune response. Such an immune response may be associated with, but is not limited to, disorders such as AIDS, Addison's disease, adult respiratory distress syndrome, allergies, anemia, asthma, atherosclerosis, bronchitis, cholecystitis, Crohn's disease, ulcerative colitis, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, erythema nodosum, atrophic gastritis, glomerulonephritis, gout, Graves' disease, hypereosinophilia, irritable bowel syndrome, lupus erythematosus, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, rheumatoid arthritis, scleroderma, Sjögren's syndrome, and autoimmune thyroiditis; complications of cancer, hemodialysis, and extracorporeal circulation; viral, bacterial, fungal, parasitic, protozoal, and helminthic infections; and trauma. In one aspect, an antibody which specifically binds NVRK1 may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express NVRK1.

In another embodiment, a vector expressing the complement of the polynucleotide encoding NVRK1 may be administered to a subject to treat or prevent an immune response as associated with, but not limited to, the disorders described above.

In another embodiment, an antagonist of NVRK1 or a fragment or derivative thereof may be administered to a subject to treat or prevent a cancer. Such cancers include, but are not limited to, adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, and teratocarcinoma; and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus. In one aspect, an antibody which specifically binds NVRK1 may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express NVRK1.

In another embodiment, a vector capable of expressing the complement of the polynucleotide encoding NVRK1, may be administered to a subject to treat or prevent a

cancer including, but not limited to, those cancers listed above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of NVRK1 may be produced using methods which are generally known in the art. In particular, purified NVRK1 may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind NVRK1.

Antibodies to NVRK1 may be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, single chain, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies, (i.e., those which inhibit dimer formation) are especially preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others, may be immunized by injection with NVRK1 or any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to NVRK1 have an amino acid sequence consisting of at least five amino acids and more preferably at least 10 amino acids. It is also preferable that they are identical to a portion of the amino acid sequence of the natural protein, and they may contain the entire amino acid sequence of a small, naturally occurring molecule. Short stretches of NVRK1 amino acids may be fused with those of another protein such as keyhole limpet hemocyanin and antibody produced against the chimeric molecule.

Monoclonal antibodies to NVRK1 may be prepared using any technique which

provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique (Kohler, G. et al. (1975) *Nature* 256:495-497; Kozbor, D. et al. (1985) *J. Immunol. Methods* 81:31-42; Cote, R.J. et al. (1983) *Proc. Natl. Acad. Sci.* 80:2026-2030; Cole, S.P. et al. (1984) *Mol. Cell Biol.* 62:109-120).

In addition, techniques developed for the production of "chimeric antibodies", the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity can be used (Morrison, S.L. et al. (1984) *Proc. Natl. Acad. Sci.* 81:6851-6855; Neuberger, M.S. et al. (1984) *Nature* 312:604-608; Takeda, S. et al. (1985) *Nature* 314:452-454). Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce NVRK1-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries (Burton D.R. (1991) *Proc. Natl. Acad. Sci.* 88:11120-3).

Antibodies may also be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature (Orlandi, R. et al. (1989) *Proc. Natl. Acad. Sci.* 86:3833-3837; Winter, G. et al. (1991) *Nature* 349:293-299).

Antibody fragments which contain specific binding sites for NVRK1 may also be generated. For example, such fragments include, but are not limited to, the F(ab')<sub>2</sub> fragments which can be produced by pepsin digestion of the antibody molecule and the Fab fragments which can be generated by reducing the disulfide bridges of the F(ab')<sub>2</sub> fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity (Huse, W.D. et al. (1989) *Science* 254:1275-1281).

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between NVRK1 and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering NVRK1 epitopes is preferred, but a competitive binding assay may also be employed (Maddox, *supra*).

In another embodiment of the invention, the polynucleotides encoding NVRK1, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding NVRK1 may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding NVRK1. Thus, complementary molecules or fragments may be used to modulate NVRK1 activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligonucleotides or larger fragments, can be designed from various locations along the coding or control regions of sequences encoding NVRK1.

Expression vectors derived from retro viruses, adenovirus, herpes or vaccinia viruses, or from various bacterial plasmids may be used for delivery of nucleotide sequences to the targeted organ, tissue or cell population. Methods which are well known to those skilled in the art can be used to construct vectors which will express nucleic acid sequence which is complementary to the polynucleotides of the gene encoding NVRK1. These techniques are described both in Sambrook et al. (supra) and in Ausubel et al. (supra).

Genes encoding NVRK1 can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide or fragment thereof which encodes NVRK1. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled by endogenous nucleases. Transient expression may last for a month or more with a non-replicating vector and even longer if appropriate replication elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing complementary sequences or antisense molecules (DNA, RNA, or PNA) to the control, 5' or regulatory regions of the gene encoding NVRK1 (signal sequence, promoters, enhancers, and introns). Oligonucleotides derived from the transcription initiation site, e.g., between positions -10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using "triple helix" base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature (Gee, J.E. et al. (1994) In: Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing Co., Mt. Kisco, NY). The complementary sequence or antisense molecule may also be designed to

block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by  
5 endonucleolytic cleavage. Examples which may be used include engineered hammerhead motif ribozyme molecules that can specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding NVRK1.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites which include the  
10 following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target gene containing the cleavage site may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease  
15 protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in  
20 vitro and in vivo transcription of DNA sequences encoding NVRK1. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA constitutively or inducibly can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life.  
25 Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-,  
30 thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced

into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections or polycationic amino polymers (Goldman, C.K. et al. (1997) Nature Biotechnology 15:462-66; incorporated herein by reference) may be achieved using methods which are well known in the art.

5 Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

An additional embodiment of the invention relates to the administration of a pharmaceutical composition, in conjunction with a pharmaceutically acceptable carrier, for  
10 any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of NVRK1, antibodies to NVRK1, mimetics, agonists, antagonists, or inhibitors of NVRK1. The compositions may be administered alone or in combination with at least one other agent, such as stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier, including, but not limited to, saline, buffered saline,  
15 dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs or hormones.

The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous,  
20 intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be  
25 found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing Co., Easton, PA).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as  
30 tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combination of active compounds with solid excipient, optionally grinding a resulting mixture, and

processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, or sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums including arabic and tragacanth; and proteins such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate.

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with a filler or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Non-lipid polycationic amino polymers may also be used for delivery. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a

manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation may be a lyophilized powder which may contain any or all of the following: 1-50 mM histidine, 0.1%-2% sucrose, and 2-7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of NVRK1, such labeling would include amount, frequency, and method of administration.

Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models, usually mice, rabbits, dogs, or pigs. The animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example NVRK1 or fragments thereof, antibodies of NVRK1, agonists, antagonists or inhibitors of NVRK1, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., ED50 (the dose therapeutically effective in 50% of the population) and LD50 (the dose lethal to 50% of the population). The dose ratio between therapeutic and toxic effects is the therapeutic index, and it can be expressed as the ratio, LD50/ED50.

Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the patient, and



the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject that requires treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or once every two weeks depending on half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from 0.1 to 100,000 micrograms, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

## DIAGNOSTICS

In another embodiment, antibodies which specifically bind NVRK1 may be used for the diagnosis of conditions or diseases characterized by expression of NVRK1, or in assays to monitor patients being treated with NVRK1, agonists, antagonists or inhibitors. The antibodies useful for diagnostic purposes may be prepared in the same manner as those described above for therapeutics. Diagnostic assays for NVRK1 include methods which utilize the antibody and a label to detect NVRK1 in human body fluids or extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by joining them, either covalently or non-covalently, with a reporter molecule. A wide variety of reporter molecules which are known in the art may be used, several of which are described above.

A variety of protocols including ELISA, RIA, and FACS for measuring NVRK1 are known in the art and provide a basis for diagnosing altered or abnormal levels of NVRK1 expression. Normal or standard values for NVRK1 expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, preferably human, with antibody to NVRK1 under conditions suitable for complex formation. The amount of standard complex formation may be quantified by various methods, but preferably by

photometric, means. Quantities of NVRK1 expressed in subject, control and disease, samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding NVRK1 may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantitate gene expression in biopsied tissues in which expression of NVRK1 may be correlated with disease. The diagnostic assay may be used to distinguish between absence, presence, and excess expression of NVRK1, and to monitor regulation of NVRK1 levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding NVRK1 or closely related molecules, may be used to identify nucleic acid sequences which encode NVRK1. The specificity of the probe, whether it is made from a highly specific region, e.g., 10 unique nucleotides in the 5' regulatory region, or a less specific region, e.g., especially in the 3' coding region, and the stringency of the hybridization or amplification (maximal, high, intermediate, or low) will determine whether the probe identifies only naturally occurring sequences encoding NVRK1, alleles, or related sequences.

Probes may also be used for the detection of related sequences, and should preferably contain at least 50% of the nucleotides from any of the NVRK1 encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and derived from the nucleotide sequence of SEQ ID NO:2 or from genomic sequence including promoter, enhancer elements, and introns of the naturally occurring NVRK1.

Means for producing specific hybridization probes for DNAs encoding NVRK1 include the cloning of nucleic acid sequences encoding NVRK1 or NVRK1 derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, radionuclides such as <sup>32</sup>P or <sup>35</sup>S, or enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding NVRK1 may be used for the diagnosis of conditions or disorders which are associated with expression of NVRK1. Examples of such

conditions or disorders include adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, and teratocarcinoma; and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; and immune responses associated with disorders such as AIDS, Addison's disease, adult respiratory distress syndrome, allergies, anemia, asthma, atherosclerosis, bronchitis, cholecystitis, Crohn's disease, ulcerative colitis, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, erythema nodosum, atrophic gastritis, glomerulonephritis, gout, Graves' disease, hypereosinophilia, irritable bowel syndrome, lupus erythematosus, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, rheumatoid arthritis, scleroderma, Sjögren's syndrome, and autoimmune thyroiditis; complications of cancer, hemodialysis, and extracorporeal circulation; viral, bacterial, fungal, parasitic, protozoal, and helminthic infections; and trauma. The polynucleotide sequences encoding NVRK1 may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; or in dipstick, pin, ELISA assays or microarrays utilizing fluids or tissues from patient biopsies to detect altered NVRK1 expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding NVRK1 may be useful in assays that detect activation or induction of various cancers, particularly those mentioned above. The nucleotide sequences encoding NVRK1 may be labeled by standard methods, and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantitated and compared with a standard value. If the amount of signal in the biopsied or extracted sample is significantly altered from that of a comparable control sample, the nucleotide sequences have hybridized with nucleotide sequences in the sample, and the presence of altered levels of nucleotide sequences encoding NVRK1 in the sample indicates the presence of the associated disease. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or in monitoring the treatment of an individual patient.

In order to provide a basis for the diagnosis of disease associated with expression of NVRK1, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either

animal or human, with a sequence, or a fragment thereof, which encodes NVRK1, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with those from an experiment where a known amount of a substantially purified polynucleotide is used.

- 5 Standard values obtained from normal samples may be compared with values obtained from samples from patients who are symptomatic for disease. Deviation between standard and subject values is used to establish the presence of disease.

Once disease is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to evaluate whether the level of expression in the patient  
10 begins to approximate that which is observed in the normal patient. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of a relatively high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the  
15 disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding  
20 NVRK1 may involve the use of PCR. Such oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably consist of two nucleotide sequences, one with sense orientation (5'→3') and another with antisense (3'←5'), employed under optimized conditions for identification of a specific gene or condition. The same two oligomers, nested sets of oligomers, or even a degenerate pool of oligomers may be  
25 employed under less stringent conditions for detection and/or quantitation of closely related DNA or RNA sequences.

Methods which may also be used to quantitate the expression of NVRK1 include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and standard curves onto which the experimental results are interpolated (Melby, P.C. et al.  
30 (1993) J. Immunol. Methods, 159:235-244; Duplaa, C. et al. (1993) Anal. Biochem. 229-236). The speed of quantitation of multiple samples may be accelerated by running the assay in an ELISA format where the oligomer of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, an oligonucleotide derived from any of the polynucleotide sequences described herein may be used as a target in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously (to produce a transcript image), and to identify genetic variants, mutations and polymorphisms. This information will be useful in determining gene function, understanding the genetic basis of disease, diagnosing disease, and in developing and monitoring the activity of therapeutic agents (Heller, R. et al. (1997) Proc. Natl. Acad. Sci. 94:2150-55) .

In one embodiment, the microarray is prepared and used according to the methods described in PCT application WO95/11995 (Chee et al.), Lockhart, D. J. et al. (1996; Nat. Biotech. 14: 1675-1680) and Schena, M. et al. (1996; Proc. Natl. Acad. Sci. 93: 10614-10619), all of which are incorporated herein in their entirety by reference.

The microarray is preferably composed of a large number of unique, single-stranded nucleic acid sequences, usually either synthetic antisense oligonucleotides or fragments of cDNAs, fixed to a solid support. The oligonucleotides are preferably about 6-60 nucleotides in length, more preferably 15-30 nucleotides in length, and most preferably about 20-25 nucleotides in length. For a certain type of microarray, it may be preferable to use oligonucleotides which are only 7-10 nucleotides in length. The microarray may contain oligonucleotides which cover the known 5', or 3', sequence, sequential oligonucleotides which cover the full length sequence; or unique oligonucleotides selected from particular areas along the length of the sequence. Polynucleotides used in the microarray may be oligonucleotides that are specific to a gene or genes of interest in which at least a fragment of the sequence is known or that are specific to one or more unidentified cDNAs which are common to a particular cell type, developmental or disease state.

In order to produce oligonucleotides to a known sequence for a microarray, the gene of interest is examined using a computer algorithm which starts at the 5' or more preferably at the 3' end of the nucleotide sequence. The algorithm identifies oligomers of defined length that are unique to the gene, have a GC content within a range suitable for hybridization, and lack predicted secondary structure that may interfere with hybridization. In certain situations it may be appropriate to use pairs of oligonucleotides on a microarray. The "pairs" will be identical, except for one nucleotide which preferably is located in the center of the sequence. The second oligonucleotide in the pair (mismatched by one) serves as a control. The number of oligonucleotide pairs may range from two to one million. The oligomers are synthesized at designated areas on a substrate using a light-directed chemical process. The substrate may be

paper, nylon or other type of membrane, filter, chip, glass slide or any other suitable solid support.

In another aspect, an oligonucleotide may be synthesized on the surface of the substrate by using a chemical coupling procedure and an ink jet application apparatus, as described in PCT application WO95/251116 (Baldeschweiler et al.) which is incorporated herein in its entirety by reference. In another aspect, a "gridded" array analogous to a dot (or slot) blot may be used to arrange and link cDNA fragments or oligonucleotides to the surface of a substrate using a vacuum system, thermal, UV, mechanical or chemical bonding procedures. An array, such as those described above, may be produced by hand or by using available devices (slot blot or dot blot apparatus), materials (any suitable solid support), and machines (including robotic instruments), and may contain 8, 24, 96, 384, 1536 or 6144 oligonucleotides, or any other number between two and one million which lends itself to the efficient use of commercially available instrumentation.

In order to conduct sample analysis using a microarray, the RNA or DNA from a biological sample is made into hybridization probes. The mRNA is isolated, and cDNA is produced and used as a template to make antisense RNA (aRNA). The aRNA is amplified in the presence of fluorescent nucleotides, and labeled probes are incubated with the microarray so that the probe sequences hybridize to complementary oligonucleotides of the microarray. Incubation conditions are adjusted so that hybridization occurs with precise complementary matches or with various degrees of less complementarity. After removal of nonhybridized probes, a scanner is used to determine the levels and patterns of fluorescence. The scanned images are examined to determine degree of complementarity and the relative abundance of each oligonucleotide sequence on the microarray. The biological samples may be obtained from any bodily fluids (such as blood, urine, saliva, phlegm, gastric juices, etc.), cultured cells, biopsies, or other tissue preparations. A detection system may be used to measure the absence, presence, and amount of hybridization for all of the distinct sequences simultaneously. This data may be used for large scale correlation studies on the sequences, mutations, variants, or polymorphisms among samples.

In another embodiment of the invention, the nucleic acid sequences which encode NVRK1 may also be used to generate hybridization probes which are useful for mapping the naturally occurring genomic sequence. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome or to artificial chromosome constructions, such as human artificial chromosomes (HACs), yeast artificial chromosomes (YACs),

bacterial artificial chromosomes (BACs), bacterial P1 constructions or single chromosome cDNA libraries as reviewed in Price, C.M. (1993) *Blood Rev.* 7:127-134, and Trask, B.J. (1991) *Trends Genet.* 7:149-154.

Fluorescent in situ hybridization (FISH as described in Verma et al. (1988) Human Chromosomes: A Manual of Basic Techniques, Pergamon Press, New York, NY) may be correlated with other physical chromosome mapping techniques and genetic map data. Examples of genetic map data can be found in various scientific journals or at Online Mendelian Inheritance in Man (OMIM). Correlation between the location of the gene encoding NVRK1 on a physical chromosomal map and a specific disease, or predisposition to a specific disease, may help delimit the region of DNA associated with that genetic disease. The nucleotide sequences of the subject invention may be used to detect differences in gene sequences between normal, carrier, or affected individuals.

In situ hybridization of chromosomal preparations and physical mapping techniques such as linkage analysis using established chromosomal markers may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms, or parts thereof, by physical mapping. This provides valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localized by genetic linkage to a particular genomic region, for example, AT to 11q22-23 (Gatti, R.A. et al. (1988) *Nature* 336:577-580), any sequences mapping to that area may represent associated or regulatory genes for further investigation. The nucleotide sequence of the subject invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc. among normal, carrier, or affected individuals.

In another embodiment of the invention, NVRK1, its catalytic or immunogenic fragments or oligopeptides thereof, can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes, between NVRK1 and the agent being tested, may be measured.

Another technique for drug screening which may be used provides for high throughput screening of compounds having suitable binding affinity to the protein of interest as described

in published PCT application WO84/03564. In this method, as applied to NVRK1 large numbers of different small test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The test compounds are reacted with NVRK1, or fragments thereof, and washed. Bound NVRK1 is then detected by methods well known in the art. Purified NVRK1 can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding NVRK1 specifically compete with a test compound for binding NVRK1. In this manner, the antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with NVRK1.

In additional embodiments, the nucleotide sequences which encode NVRK1 may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

The examples below are provided to illustrate the subject invention and are not included for the purpose of limiting the invention.

## EXAMPLES

### I PROSTUT04 cDNA Library Construction

The PROSTUT04 cDNA library was constructed from prostate tumor tissue obtained from a 57-year-old Caucasian male. Surgery included a radical prostatectomy, removal of both testes and excision of regional lymph nodes. The pathology report indicated an adenocarcinoma (Gleason grade 3+3) in both the left and right periphery of the prostate.

Perineural invasion was present, as was involvement of periprostatic tissue. A single right pelvic lymph node, the right and left apical surgical margins were positive for tumor; the seminal vesicles were negative. The patient history includes a benign neoplasm of the large bowel, appendectomy, and tonsillectomy with adenoidectomy. The patient was taking insulin for type I diabetes. The patient's family history included a malignant neoplasm of the prostate in the patient's father and type I diabetes without complications in the mother.

The frozen tissues were homogenized and lysed using a Brinkmann Homogenizer Polytron PT-3000 (Brinkmann Instruments, Westbury, NJ) in guanidinium isothiocyanate solution. Both lysates were extracted once with an equal volume acid phenol per Stratgene's



RNA isolation protocol (Stratagene Inc., San Diego, CA). The RNA was extracted once with acid phenol pH 4.7, precipitated using 0.3 M sodium acetate and 2.5 volumes of ethanol, resuspended in DEPC-treated water, and DNase treated at 37°C for 25 minutes. The mRNA was then isolated using the Qiagen Oligotex kit (QIAGEN, Inc., Chatsworth, CA) and used to  
5 construct the cDNA libraries. cDNAs were fractionated on a Sepharose CL4B column (Cat. #275105-01, Pharmacia), and those cDNAs exceeding 400 bp were ligated into pSport I. The plasmid pSport I was subsequently transformed into DH5a™ competent cells (Cat. #18258-012, Gibco/BRL).

The RNA was handled according to the recommended protocols in the SuperScript  
10 Plasmid System for cDNA Synthesis and Plasmid Cloning (Cat. #18248-013; Gibco/BRL). cDNAs were fractionated on a Sepharose CL4B column (Cat. 275105, Pharmacia), and those cDNAs exceeding 400 bp were ligated into pSport I. The plasmid pSport I was subsequently transformed into DH5a™ competent cells (Cat. #18258-012, Gibco/BRL).

## 15 II Isolation and Sequencing of cDNA Clones

Plasmid DNA was released from the cells and purified using the REAL Prep 96 Plasmid Kit (Catalog #26173, QIAGEN, Inc.). This kit enabled the simultaneous purification of 96 samples in a 96-well block using multi-channel reagent dispensers. The recommended protocol was employed except for the following changes: 1) the bacteria were cultured in 1  
20 ml of sterile Terrific Broth (Catalog #22711, GIBCO/BRL) with carbenicillin at 25 mg/L and glycerol at 0.4%; 2) after inoculation, the cultures were incubated for 19 hours and at the end of incubation, the cells were lysed with 0.3 ml of lysis buffer; and 3) following isopropanol precipitation, the plasmid DNA pellet was resuspended in 0.1 ml of distilled water. After the last step in the protocol, samples were transferred to a 96-well block for storage at 4° C.

25 The cDNAs were sequenced by the method of Sanger et al. (1975, J. Mol. Biol. 94:441f), using a Hamilton Micro Lab 2200 (Hamilton, Reno, NV) in combination with Peltier Thermal Cyclers (PTC200 from MJ Research, Watertown, MA) and Applied Biosystems 377 DNA Sequencing Systems; and the reading frame was determined.

## 30 III Homology Searching of cDNA Clones and Their Deduced Proteins

The nucleotide sequences of the Sequence Listing or amino acid sequences deduced from them were used as query sequences against databases such as GenBank, SwissProt, BLOCKS, and Pima II. These databases which contain previously identified and annotated

sequences were searched for regions of homology (similarity) using BLAST, which stands for Basic Local Alignment Search Tool (Altschul (1993) supra, Altschul (1990) supra).

BLAST produces alignments of both nucleotide and amino acid sequences to determine sequence similarity. Because of the local nature of the alignments, BLAST is especially useful in determining exact matches or in identifying homologs which may be of prokaryotic (bacterial) or eukaryotic (animal, fungal or plant) origin. Other algorithms such as the one described in Smith et al. (1992 Protein Engineering 5:35-51), incorporated herein by reference, can be used when dealing with primary sequence patterns and secondary structure gap penalties. As disclosed in this application, the sequences have lengths of at least 49 nucleotides, and no more than 12% uncalled bases (where N is recorded rather than A, C, G, or T).

The BLAST approach, as detailed in Karlin and Altschul (supra) and incorporated herein by reference, searches for matches between a query sequence and a database sequence, to evaluate the statistical significance of any matches found, and to report only those matches which satisfy the user-selected threshold of significance. In this application, threshold was set at  $10^{-25}$  for nucleotides and  $10^{-14}$  for peptides.

Incyte nucleotide sequences were searched against the GenBank databases for primate (pri), rodent (rod), and mammalian sequences (mam), and deduced amino acid sequences from the same clones are searched against GenBank functional protein databases, mammalian (mamp), vertebrate (vrtp) and eukaryote (eukp), for homology.

#### IV Northern Analysis

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound (Sambrook et al., supra).

Analogous computer techniques using BLAST (Altschul, S.F. (1993) J.Mol.Evol. 36:290-300; Altschul, S.F. et al. (1990) J.Mol.Evol. 215:403-410) are used to search for identical or related molecules in nucleotide databases such as GenBank or the LIFESEQ™ database (Incyte Pharmaceuticals). This analysis is much faster than multiple, membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or homologous.

The basis of the search is the product score which is defined as:

$$\underline{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1-2% error; and at 70, the match will be exact. Homologous molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

The results of northern analysis are reported as a list of libraries in which the transcript encoding NVRK1 occurs. Abundance and percent abundance are also reported. Abundance directly reflects the number of times a particular transcript is represented in a cDNA library, and percent abundance is abundance divided by the total number of sequences examined in the cDNA library.

## V Extension of NVRK1 Encoding Polynucleotides

The nucleic acid sequence of the Incyte Clone 1543990 was used to design oligonucleotide primers for extending a partial nucleotide sequence to full length. One primer was synthesized to initiate extension in the antisense direction, and the other was synthesized to extend sequence in the sense direction. Primers were used to facilitate the extension of the known sequence "outward" generating amplicons containing new, unknown nucleotide sequence for the region of interest. The initial primers were designed from the cDNA using OLIGO 4.06 (National Biosciences), or another appropriate program, to be about 22 to about 30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures of about 68° to about 72° C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries (Gibco/BRL) were used to extend the sequence. If more than one extension is necessary or desired, additional sets of primers are designed to further extend the known region.

High fidelity amplification was obtained by following the instructions for the XL-PCR kit (Perkin Elmer) and thoroughly mixing the enzyme and reaction mix. Beginning with 40 pmol of each primer and the recommended concentrations of all other components of the kit, PCR was performed using the Peltier Thermal Cycler (PTC200; M.J. Research, Watertown, MA) and the following parameters:

Step 1	94° C for 1 min (initial denaturation)
Step 2	65° C for 1 min
Step 3	68° C for 6 min

	Step 4	94° C for 15 sec
	Step 5	65° C for 1 min
	Step 6	68° C for 7 min
	Step 7	Repeat step 4-6 for 15 additional cycles
5	Step 8	94° C for 15 sec
	Step 9	65° C for 1 min
	Step 10	68° C for 7:15 min
	Step 11	Repeat step 8-10 for 12 cycles
	Step 12	72° C for 8 min
10	Step 13	4° C (and holding)

A 5-10  $\mu$ l aliquot of the reaction mixture was analyzed by electrophoresis on a low concentration (about 0.6-0.8%) agarose mini-gel to determine which reactions were successful in extending the sequence. Bands thought to contain the largest products were excised from the gel, purified using QIAQuick™ (QIAGEN Inc., Chatsworth, CA), and trimmed of overhangs using Klenow enzyme to facilitate religation and cloning.

After ethanol precipitation, the products were redissolved in 13  $\mu$ l of ligation buffer, 1  $\mu$ l T4-DNA ligase (15 units) and 1  $\mu$ l T4 polynucleotide kinase were added, and the mixture was incubated at room temperature for 2-3 hours or overnight at 16° C. Competent *E. coli* cells (in 40  $\mu$ l of appropriate media) were transformed with 3  $\mu$ l of ligation mixture and cultured in 80  $\mu$ l of SOC medium (Sambrook et al., supra). After incubation for one hour at 37° C, the *E. coli* mixture was plated on Luria Bertani (LB)-agar (Sambrook et al., supra) containing 2x Carb. The following day, several colonies were randomly picked from each plate and cultured in 150  $\mu$ l of liquid LB/2x Carb medium placed in an individual well of an appropriate, commercially-available, sterile 96-well microtiter plate. The following day, 5  $\mu$ l of each overnight culture was transferred into a non-sterile 96-well plate and after dilution 1:10 with water, 5  $\mu$ l of each sample was transferred into a PCR array.

For PCR amplification, 18  $\mu$ l of concentrated PCR reaction mix (3.3x) containing 4 units of rTth DNA polymerase, a vector primer, and one or both of the gene specific primers used for the extension reaction were added to each well. Amplification was performed using the following conditions:

	Step 1	94° C for 60 sec
	Step 2	94° C for 20 sec
	Step 3	55° C for 30 sec
35	Step 4	72° C for 90 sec
	Step 5	Repeat steps 2-4 for an additional 29 cycles
	Step 6	72° C for 180 sec
	Step 7	4° C (and holding)

40 Aliquots of the PCR reactions were run on agarose gels together with molecular

weight markers. The sizes of the PCR products were compared to the original partial cDNAs, and appropriate clones were selected, ligated into plasmid, and sequenced.

In like manner, the nucleotide sequence of SEQ ID NO:2 is used to obtain 5' regulatory sequences using the procedure above, oligonucleotides designed for 5' extension, and an appropriate genomic library.

## VI Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:2 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base-pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 (National Biosciences), labeled by combining 50 pmol of each oligomer and 250  $\mu$ Ci of [ $\gamma$ - $^{32}$ P] adenosine triphosphate (Amersham) and T4 polynucleotide kinase (DuPont NEN<sup>®</sup>, Boston, MA). The labeled oligonucleotides are substantially purified with Sephadex G-25 superfine resin column (Pharmacia & Upjohn). A aliquot containing  $10^7$  counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases (Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II; DuPont NEN<sup>®</sup>).

The DNA from each digest is fractionated on a 0.7 percent agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham, NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under increasingly stringent conditions up to 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. After XOMAT AR<sup>™</sup> film (Kodak, Rochester, NY) is exposed to the blots in a Phosphorimager cassette (Molecular Dynamics, Sunnyvale, CA) for several hours, hybridization patterns are compared visually.

## VII Microarrays

To produce oligonucleotides for a microarray, the nucleotide sequence described herein is examined using a computer algorithm which starts at the 3' end of the nucleotide sequence. The algorithm identifies oligomers of defined length that are unique to the gene, have a GC content within a range suitable for hybridization, and lack predicted secondary structure that would interfere with hybridization. The algorithm identifies 20 sequence-specific oligonucleotides of 20 nucleotides in length (20-mers). A matched set of

oligonucleotides is created in which one nucleotide in the center of each sequence is altered. This process is repeated for each gene in the microarray, and double sets of twenty 20 mers are synthesized and arranged on the surface of the silicon chip using a light-directed chemical process (Chee, M. et al., PCT/WO95/11995, incorporated herein by reference).

5 In the alternative, a chemical coupling procedure and an ink jet device are used to synthesize oligomers on the surface of a substrate (Baldeschweiler, J.D. et al., PCT/WO95/25116, incorporated herein by reference). In another alternative, a "gridded" array analogous to a dot (or slot) blot is used to arrange and link cDNA fragments or oligonucleotides to the surface of a substrate using a vacuum system, thermal, UV,  
10 mechanical or chemical bonding procedures. An array may be produced by hand or using available materials and machines and contain grids of 8 dots, 24 dots, 96 dots, 384 dots, 1536 dots or 6144 dots. After hybridization, the microarray is washed to remove nonhybridized probes, and a scanner is used to determine the levels and patterns of fluorescence. The scanned images are examined to determine degree of complementarity and the relative  
15 abundance of each oligonucleotide sequence on the micro-array.

### VIII Complementary Polynucleotides

Sequence complementary to the NVRK1-encoding sequence, or any part thereof, is used to decrease or inhibit expression of naturally occurring NVRK1. Although use of  
20 oligonucleotides comprising from about 15 to about 30 base-pairs is described, essentially the same procedure is used with smaller or larger sequence fragments. Appropriate oligonucleotides are designed using Oligo 4.06 software and the coding sequence of NVRK1, SEQ ID NO:1. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence.  
25 To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the NVRK1-encoding transcript.

### IX Expression of NVRK1

Expression of NVRK1 is accomplished by subcloning the cDNAs into appropriate  
30 vectors and transforming the vectors into host cells. In this case, the cloning vector is also used to express NVRK1 in *E. coli*. Upstream of the cloning site, this vector contains a promoter for  $\beta$ -galactosidase, followed by sequence containing the amino-terminal Met, and the subsequent seven residues of  $\beta$ -galactosidase. Immediately following these eight residues

is a bacteriophage promoter useful for transcription and a linker containing a number of unique restriction sites.

Induction of an isolated, transformed bacterial strain with IPTG using standard methods produces a fusion protein which consists of the first eight residues of  $\beta$ -galactosidase, about 5 to 15 residues of linker, and the full length protein. The signal residues direct the secretion of NVRK1 into the bacterial growth media which can be used directly in the following assay for activity.

## **X Demonstration of NVRK1 Activity**

Samples were fractionated on SDS-polyacrylamide gel (10% (w/v) acrylamide) containing GST-LT- $\beta$ R(CD) fusion protein. The gel was then washed with 20% (v/v) propan-2-ol to remove SDS. The gel was denatured with 6 M guanidine HCl, then renatured in 5 mM 2-mercaptoethanol, 50 mM Tris-HCl, pH 8.0, and 0.04% (v/v) Tween 40. Renaturation was followed by washing at 4 °C for 30 min with two changes of buffer (20 mM HEPES, pH 7.4, 0.2 mM NaF, 0.1 mM sodium orthovanadate), and incubated at 22 °C for 3 h in the kinase reaction buffer containing 50  $\mu$ Ci of [ $\gamma$ - $^{32}$ P]ATP. After removal of unreacted [ $\gamma$ - $^{32}$ P]ATP, protein kinases were visualized by autoradiography of the dried gel. Signals due to protein kinases were also detected and quantified using a PhosphorImager<sup>TM</sup> (Molecular Dynamics) to allow comparison of signal strengths (Mei-Yi et al. (1997) J. Biol. Chem. 272 (27):17154-17159).

## **XI Production of NVRK1 Specific Antibodies**

NVRK1 that is substantially purified using PAGE electrophoresis (Sambrook, supra), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols. The amino acid sequence deduced from SEQ ID NO:2 is analyzed using DNASTAR software (DNASTAR Inc) to determine regions of high immunogenicity and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions, is described by Ausubel et al. (supra), and others.

Typically, the oligopeptides are 15 residues in length, synthesized using an Applied Biosystems Peptide Synthesizer Model 431A using fmoc-chemistry, and coupled to keyhole limpet hemocyanin (KLH, Sigma, St. Louis, MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS; Ausubel et al., supra). Rabbits are immunized with the

oligopeptide-KLH complex in complete Freund's adjuvant. The resulting antisera are tested for antipeptide activity, for example, by binding the peptide to plastic, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio iodinated, goat anti-rabbit IgG.

5

## **XII Purification of Naturally Occurring NVRK1 Using Specific Antibodies**

Naturally occurring or recombinant NVRK1 is substantially purified by immunoaffinity chromatography using antibodies specific for NVRK1. An immunoaffinity column is constructed by covalently coupling NVRK1 antibody to an activated  
10 chromatographic resin, such as CNBr-activated Sepharose (Pharmacia & Upjohn). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing NVRK1 is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of NVRK1 (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under  
15 conditions that disrupt antibody/NVRK1 binding (eg, a buffer of pH 2-3 or a high concentration of a chaotrope, such as urea or thiocyanate ion), and NVRK1 is collected.

## **XIII Identification of Molecules Which Interact with NVRK1**

NVRK1 or biologically active fragments thereof are labeled with <sup>125</sup>I Bolton-Hunter reagent (Bolton et al. (1973) Biochem. J. 133: 529). Candidate molecules previously arrayed  
20 in the wells of a multi-well plate are incubated with the labeled NVRK1, washed and any wells with labeled NVRK1 complex are assayed. Data obtained using different concentrations of NVRK1 are used to calculate values for the number, affinity, and association of NVRK1 with the candidate molecules.

25 All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed  
30 should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.



What is claimed is:

1. A substantially purified VRK1 kinase comprising the amino acid sequence of SEQ ID NO:1 or fragments thereof.
2. A substantially purified variant of the VRK1 kinase having at least 90% amino acid identity to SEQ ID NO:1 and which retains at least one functional characteristic of VRK1 kinase.
3. An isolated and purified polynucleotide sequence encoding the VRK1 kinase of claim 1 or fragments or variants of said polynucleotide sequence.
4. A composition comprising the polynucleotide sequence of claim 3.
5. A polynucleotide sequence which hybridizes to the polynucleotide sequence of claim 3.
6. A polynucleotide sequence which is complementary to the polynucleotide sequence of claim 3 or fragments or variants thereof.
7. An isolated and purified polynucleotide sequence comprising SEQ ID NO:2 or fragments or variants thereof.
8. A polynucleotide sequence which is complementary to the polynucleotide sequence of claim 7.
9. An expression vector containing at least a fragment of the polynucleotide sequence of claim 3.
10. A host cell containing the vector of claim 9.
11. A method for producing a polypeptide comprising the amino acid sequence of SEQ ID NO:1, or a fragment thereof, the method comprising the steps of:

- a) culturing the host cell of claim 10 under conditions suitable for the expression of the polypeptide; and
- b) recovering the polypeptide from the host cell culture.

5           12.    A pharmaceutical composition comprising a substantially purified VRK1 kinase having the amino acid sequence of SEQ ID NO:1 in conjunction with a suitable pharmaceutical carrier.

10           13.    A purified antibody which specifically binds to the polypeptide of claim 1.

          14.    A purified agonist of the polypeptide of claim 1.

          15.    A purified antagonist of the polypeptide of claim 1.

15           16.    A method for treating an immune response comprising administering to a subject in need of such treatment an effective amount of the antagonist of claim 15.

          17.    A method for treating a cancer comprising administering to a subject in need of such treatment an effective amount of the antagonist of claim 15.

20           18.    A method for detecting a polynucleotide which encodes the VRK1 kinase in a biological sample comprising the steps of:

          a)     hybridizing the polynucleotide of claim 6 to nucleic acid material of a biological sample, thereby forming a hybridization complex; and

25           b)     detecting said hybridization complex, wherein the presence of said complex correlates with the presence of a polynucleotide encoding VRK1 kinase in said biological sample.

30           19.    The method of claim 18 wherein the nucleic acid material is amplified by the polymerase chain reaction prior to hybridization.

1/10

9	18	27	36	45	54
NGA CCC AGG GAG CAG AAG GAC GGG GAT CTC CCT TTG CCG CTA AGA GAC TCC AAC					
63	72	81	90	99	108
TCC CAG GAT GCC TTG GGA GCG CTG GAC CAC AGC TTC CCG GCA TTT CCC CGN GNG					
117	126	135	144	153	162
GAG GGG GCT AGC GTA GCC TCG CNC CGC GCC TCG TCG TAC CAC TCG TCC CGC CCC					
171	180	189	198	207	216
TCC TCG TCG CCT CCG AGC CAA TGG GAA GGC TCC ATA CTG CAG GGT GCG AAG GGG					
225	234	243	252	261	270
CCG GCG CCG CTG CCG AGT TAC GAG TCG GCG AAA GCG GCG GGA AGT TCG TAC TGG					
279	288	297	306	315	324
GCA GAA CGC GAC GGG TCT GCG GCT TAG GTG AAA ATG CCT CGT GTA AAA GCA GCT					
			M	P	R
				V	K
					A
333	342	351	360	369	378
CAA GCT GGA AGA CAG AGC TCT GCA AAG AGA CAT CTT GCA GAA CAA TTT GCA GTT					
Q	A	G	R	Q	S
		S	S	A	K
				R	H
				L	A
				E	F
				Q	A
					V

FIGURE 1A

2/10

387	GGA GAG ATA ATA ACT GAC ATG GCA AAA AAG GAA TGG AAA GTA GGA TTA CCC ATT	414	423	432
	G E I I T D M A K K E W K V G L P I			
396		405		
441	GGC CAA GGA GGC TTT GGC TGT ATA TAT CTT GCT GAT ATG AAT TCT TCA GAG TCA	459	468	477
	G Q G G F G C I Y L A D M N S E S			
450		495	504	513
495	GTT GGC AGT GAT GCA CCT TGT GTT GTA AAA GTG GAA CCC AGT GAC AAT GGA CCT	522	531	540
	V G S D A P C V C V K V E P S D N G P			
549	CTT TTT ACT GAA TTA AAG TTC TAC CAA CGA GCT GCA AAA CCA GAG CAA ATT CAG	567	576	585
	L F T E L K F Y Q R A A K P E Q I Q			
603	AAA TGG ATT CGT ACC CGT AAG CTG AAG TAC CTG GGT GTT CCT AAG TAT TGG GGG	612	621	630
	K W I R T R K L K Y L G V P K Y W G			
657	TCT GGT CTA CAT GAC AAA AAT GGA AAA AGT TAC AGG TTT ATG ATA ATG GAT CGC	666	675	684
	S G L H D K N G K S Y R F M I M D R			
711	TTT GGG AGT GAC CTT CAG AAA ATA TAT GAA GCA AAT GCC AAA AGG TTT TCT CGG	720	729	738
	F G S D L Q K I Y E A N A K R F S R			

FIGURE 1B

3/10

765	AAA ACT GTC TTG CAG CTA AGC TTA AGA ATT CTG GAT ATT CTG GAA TAT ATT CAC	810
K T V L Q L S L R I L D I L E Y I H		
819	GAG CAT GAG TAT GTG CAT GGA GAT ATC AAG GCC TCA AAT CTT CTG AAC TAC	864
E H E Y V H G D I K A S N L L N Y		
873	AAG AAT CCT GAC CAG GTG TAC TTG GTA GAT TAT GGC CTT GCT TAT CGG TAC TGC	918
K N P D Q V Y L V D Y G L A Y R Y C		
927	CCA GAA GGA GTT CAT AAA GAA TAC AAA GAA GAC CCC AAA AGA TGT CAC GAT GGC	972
P E G V H K E Y K E D P K R C H D G		
981	ACT ATT GAA TTC ACG AGC ATC GAT GCA CAC AAT GGT GTG GCC CCA TCA AGA CGT	1026
T I E F T S I D A H N G V A P S R R		
1035	GGT GAT TTG GAA ATA CTT GGT TAT TGC ATG ATC CAA TGG CTT ACT GGC CAT CTT	1080
G D L E I L G Y C M I Q W L T G H L		
1089	CCT TGG GAG GAT AAT TTG AAA GAT CCT AAA TAT GTT AGA GAT TCC AAA ATT AGA	1134
P W E D N L K D P K Y V R D S K I R		

FIGURE 1C

4/10

1143	1152	1161	1170	1179	1188
TAC AGA GAA AAT ATT GCA AGT TTG ATG GAC AAA TGT TTT CCT GAG AAA AAC AAA					
Y R E N I A A S L M D K C F P E K N K					
1197	1206	1215	1224	1233	1242
CCA GGT GAA ATT GCC AAA TAC ATG GAA ACA GTG AAA TTA CTA GAC TAC ACT GAA					
P G E I A K Y M E T V K L L D Y T E					
1251	1260	1269	1278	1287	1296
AAA CCT CTT TAT GAA AAT TTA CGT GAC ATT CTT TTG CAA GGA CTA AAA GCT ATA					
K P L Y E N L R D I L L Q G L K A I					
1305	1314	1323	1332	1341	1350
GGA AGT AAG GAT GAT GGC AAA TTG GAC CTC AGT GTT GTG GAG AAT GGA GGT TTG					
G S K D D G K K L D L S V V E N G G L					
1359	1368	1377	1386	1395	1404
AAA GCA AAA ACA ATA ACA AAG AAG CGA AAG AAA GAA ATT GAA GAA AGC AAG GAA					
K A K T I T K K R K K E I E E S K E					
1413	1422	1431	1440	1449	1458
CCT GGT GTT GAA GAT ACG GAA TGG TCA AAC ACA CAG ACA GAG GAG GCC ATA CAG					
P G V E D T E W S N T Q T E E A I Q					
1467	1476	1485	1494	1503	1512
ACC CGT TCA AGA ACC AGA AAG AGA GTC CAG AAG TAA TTC AGA TGC TGT GAA CCA					
T R S R R T R K R V Q K					

FIGURE 1D

5/10

1521 1530 1539 1548 1557 1566  
GAT TTC CTT TTC TTT GTT TTC TTT TGA CTT TTT TCT CCT TTT CTA TTT GAA CTG

1575 1584 1593 1602 1611 1620  
TTT TAT TTT CCT GTG AGT CTT GCG AGG TGG AAG TAA TGA TTA AAT ACT CAT GTG

1629 1638 1647 1656 1665 1674  
TTC AGA AAA CAT AAA CTT TTT TTA TAA AAA TAT TTT GTA CAA TTC ATT AAA GGC

1683 1692 1701 1710 1719 1728  
TAA TTT ATG AAA TTT GAA AAT CTT CAG GTT ATA CTC CTT AAG TTA TCC CAA AGC

1737 1746 1755 1764 1773 1782  
CGT GTG TTT GTG ATG TTT TGG AGT ACA TAT ATA TGA AAA TTA TTA TGA CAC GCA

1791 1800 1809 1818 1827 1836  
CTT TTC TAA TCA TTG TAC ATT TCT CAG AGT GGA TAA AAA TGT TTG ACA AAG TCC

1845 1854 1863 1872 1881  
TCA CTT TTA AGG AAA TGC AAA GCT TAA AAT AAA ACT CTC TTT TGT TTG ATG C

FIGURE 1E

6/10

1	M	P	R	V	K	A	A	Q	A	G	R	Q	S	S	A	K	R	H	L	A	E	Q	F	A	V	G	E	I	I	T	1543990
1	M	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	N	F	Q	-	G	L	V	L	T	GI 222701	
31	D	M	A	K	K	E	W	K	V	G	L	P	I	G	Q	G	G	F	G	C	I	Y	L	A	D	M	N	S	S	E	1543990
10	D	N	C	K	N	Q	W	V	V	G	P	L	I	G	K	G	G	F	G	S	I	Y	-	-	-	-	-	-	-	-	GI 222701
61	S	V	G	S	D	A	P	C	V	V	K	V	E	P	S	D	N	G	P	L	F	T	E	L	K	F	Y	Q	R	A	1543990
32	-	T	T	N	D	N	N	Y	V	V	K	I	E	P	K	A	N	G	S	L	F	T	E	Q	A	F	Y	T	R	V	GI 222701
91	A	K	P	E	Q	I	Q	K	W	I	R	T	R	K	L	K	Y	L	G	V	P	K	Y	W	G	S	G	L	H	D	1543990
61	L	K	P	S	V	I	E	E	W	K	K	S	H	N	I	K	H	V	G	L	I	T	C	K	A	F	G	L	Y	K	GI 222701
121	K	N	G	K	S	Y	R	F	M	I	M	D	R	F	G	S	D	L	Q	K	I	Y	E	A	N	A	K	R	F	S	1543990
91	S	I	N	V	E	Y	R	F	L	V	I	N	R	L	G	A	D	L	D	A	V	I	R	A	N	N	R	L	P	GI 222701	
151	R	K	T	V	L	Q	L	S	L	R	I	L	D	I	L	E	Y	I	H	E	H	E	Y	V	H	G	D	I	K	A	1543990
121	K	R	S	V	M	L	I	G	I	E	I	L	N	T	I	Q	F	M	H	E	Q	G	Y	S	H	G	D	I	K	A	GI 222701
181	S	N	L	L	N	Y	K	N	P	D	Q	V	Y	L	V	D	Y	G	L	A	Y	R	Y	C	P	E	G	V	H	1543990	
151	S	N	I	V	L	D	Q	I	D	K	N	K	L	Y	L	V	D	Y	G	L	V	S	K	F	M	S	N	G	E	H	GI 222701
211	K	E	Y	K	E	D	P	K	R	C	H	D	G	T	I	E	F	T	S	I	D	A	H	N	G	V	A	P	S	R	1543990
181	V	P	F	I	R	N	P	N	K	M	D	N	G	T	L	E	F	T	P	I	D	S	H	K	G	Y	V	V	S	R	GI 222701

FIGURE 2A



7/10

241	R	G	D	L	E	I	L	G	Y	C	M	I	Q	W	L	T	G	H	L	P	W	E	D	-	-	N	L	K	D	P	1543990
211	R	G	D	L	E	T	L	G	Y	C	M	I	R	W	L	G	G	I	L	P	W	T	K	I	S	E	T	K	N	C	GI 222701
269	K	Y	V	R	D	S	K	I	R	Y	R	E	N	I	A	S	L	M	D	K	C	F	P	E	K	N	K	P	G	E	1543990
241	A	L	V	S	A	T	K	Q	K	Y	V	N	N	T	A	T	L	L	M	T	S	L	-	-	Q	Y	A	P	R	E	GI 222701
299	I	A	K	Y	M	E	T	V	K	L	L	D	Y	T	E	K	P	L	Y	E	N	L	R	D	I	L	L	Q	G	L	1543990
269	L	L	Q	Y	I	T	M	V	N	S	L	T	Y	F	E	E	P	N	Y	D	E	F	R	H	I	L	M	Q	G	-	GI 222701
329	K	A	I	G	S	K	D	D	G	K	L	D	L	S	V	V	E	N	G	G	L	K	A	K	T	I	T	K	K	R	1543990
298	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	GI 222701
359	K	K	E	I	E	E	S	K	E	P	G	V	E	D	T	E	W	S	N	T	Q	T	E	E	A	I	Q	T	R	S	1543990
298	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	GI 222701
389	R	T	R	K	R	V	Q	K																							1543990
298	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	GI 222701

FIGURE 2B

8/10




FIGURE 3B

9/10




FIGURE 3A

10/10




FIGURE 4

## SEQUENCE LISTING

&lt;110&gt; INCYTE PHARMACEUTICALS, INC.

YUE, Henry  
LAL, Preeti  
CORLEY, Neil c.

&lt;120&gt; NEW VRK1 KINASE

&lt;130&gt; PF-0378 PCT

&lt;140&gt; To Be Assigned

&lt;141&gt; Herewith

&lt;150&gt; US 08/923,469

&lt;151&gt; 1997-09-01

&lt;160&gt; 3

&lt;170&gt; PERL Program

&lt;210&gt; 1

&lt;211&gt; 396

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt; -

&lt;223&gt; 1543990

&lt;400&gt; 1

Met	Pro	Arg	Val	Lys	Ala	Ala	Gln	Ala	Gly	Arg	Gln	Ser	Ser	Ala
1				5					10					15
Lys	Arg	His	Leu	Ala	Glu	Gln	Phe	Ala	Val	Gly	Glu	Ile	Ile	Thr
				20					25					30
Asp	Met	Ala	Lys	Lys	Glu	Trp	Lys	Val	Gly	Leu	Pro	Ile	Gly	Gln
				35					40					45
Gly	Gly	Phe	Gly	Cys	Ile	Tyr	Leu	Ala	Asp	Met	Asn	Ser	Ser	Glu
				50					55					60
Ser	Val	Gly	Ser	Asp	Ala	Pro	Cys	Val	Val	Lys	Val	Glu	Pro	Ser
				65					70					75
Asp	Asn	Gly	Pro	Leu	Phe	Thr	Glu	Leu	Lys	Phe	Tyr	Gln	Arg	Ala
				80					85					90
Ala	Lys	Pro	Glu	Gln	Ile	Gln	Lys	Trp	Ile	Arg	Thr	Arg	Lys	Leu
				95					100					105
Lys	Tyr	Leu	Gly	Val	Pro	Lys	Tyr	Trp	Gly	Ser	Gly	Leu	His	Asp
				110					115					120
Lys	Asn	Gly	Lys	Ser	Tyr	Arg	Phe	Met	Ile	Met	Asp	Arg	Phe	Gly
				125					130					135
Ser	Asp	Leu	Gln	Lys	Ile	Tyr	Glu	Ala	Asn	Ala	Lys	Arg	Phe	Ser
				140					145					150
Arg	Lys	Thr	Val	Leu	Gln	Leu	Ser	Leu	Arg	Ile	Leu	Asp	Ile	Leu
				155					160					165
Glu	Tyr	Ile	His	Glu	His	Glu	Tyr	Val	His	Gly	Asp	Ile	Lys	Ala
				170					175					180
Ser	Asn	Leu	Leu	Leu	Asn	Tyr	Lys	Asn	Pro	Asp	Gln	Val	Tyr	Leu
				185					190					195
Val	Asp	Tyr	Gly	Leu	Ala	Tyr	Arg	Tyr	Cys	Pro	Glu	Gly	Val	His
				200					205					210
Lys	Glu	Tyr	Lys	Glu	Asp	Pro	Lys	Arg	Cys	His	Asp	Gly	Thr	Ile
				215					220					225
Glu	Phe	Thr	Ser	Ile	Asp	Ala	His	Asn	Gly	Val	Ala	Pro	Ser	Arg
				230					235					240
Arg	Gly	Asp	Leu	Glu	Ile	Leu	Gly	Tyr	Cys	Met	Ile	Gln	Trp	Leu
				245					250					255
Thr	Gly	His	Leu	Pro	Trp	Glu	Asp	Asn	Leu	Lys	Asp	Pro	Lys	Tyr
				260					265					270

Val	Arg	Asp	Ser	Lys	Ile	Arg	Tyr	Arg	Glu	Asn	Ile	Ala	Ser	Leu
				275					280					285
Met	Asp	Lys	Cys	Phe	Pro	Glu	Lys	Asn	Lys	Pro	Gly	Glu	Ile	Ala
				290					295					300
Lys	Tyr	Met	Glu	Thr	Val	Lys	Leu	Leu	Asp	Tyr	Thr	Glu	Lys	Pro
				305					310					315
Leu	Tyr	Glu	Asn	Leu	Arg	Asp	Ile	Leu	Leu	Gln	Gly	Leu	Lys	Ala
				320					325					330
Ile	Gly	Ser	Lys	Asp	Asp	Gly	Lys	Leu	Asp	Leu	Ser	Val	Val	Glu
				335					340					345
Asn	Gly	Gly	Leu	Lys	Ala	Lys	Thr	Ile	Thr	Lys	Lys	Arg	Lys	Lys
				350					355					360
Glu	Ile	Glu	Glu	Ser	Lys	Glu	Pro	Gly	Val	Glu	Asp	Thr	Glu	Trp
				365					370					375
Ser	Asn	Thr	Gln	Thr	Glu	Glu	Ala	Ile	Gln	Thr	Arg	Ser	Arg	Thr
				380					385					390
Arg	Lys	Arg	Val	Gln	Lys									
				395										

&lt;210&gt; 2

&lt;211&gt; 1887

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; 104, 106, 130

&lt;223&gt; a or g or c or t, unknown, or other

&lt;220&gt; -

&lt;223&gt; 1543990

&lt;400&gt; 2

gacccaggga	gcagaaggac	ggggatctcc	ctttgccgct	aagagactcc	aactcccagg	60
atgccttgga	agcgctggac	cacagcttcc	cggcatttcc	ccgngnggag	ggggctagcg	120
tagcctcgcn	ccgcgcctcg	cgttaccact	cgtcccgcgc	ctcctcgteg	cctccgagcc	180
aatgggaagg	ctccatactg	cagggtgcga	aggggccggc	gccgctgccc	agttacgagt	240
cggcgaaagc	ggcggaaggt	tcgtactggg	cagaacgcga	cgggtctgcg	gcttaggtga	300
aaatgcctcg	tgtaaaagca	gctcaagctg	gaagacagag	ctctgcaaag	agacatcttg	360
cagaacaatt	tgcagttgga	gagataataa	ctgacatggc	aaaaaaggaa	tggaaagtag	420
gattaccocat	tggccaagga	ggccttgggt	gtatatactt	tgctgatatg	aattcttcag	480
agtcagttgg	cagtgatgca	ccttggtgtg	taaaagtggg	acccagtgac	aatggacctc	540
tttttactga	attaaagtgc	taccaacgag	ctgcaaaacc	agagcaaatt	cagaaatgga	600
ttcgtaccgc	taagctgaag	tacctgggtg	ttcctaagta	ttgggggtct	ggtctacatg	660
acaaaaatgg	aaaaagttac	aggtttatga	taatggatcg	ctttgggagt	gaccttcaga	720
aaatatatga	agcaaagtc	aaaaggtttt	ctcggaatac	tgtcttgagc	ctaagcttaa	780
gaattcttga	tattctggaa	tattttcacg	agcatgagta	tgtgcatgga	gatatcaagg	840
cctcaaatct	tcttctgaac	tacaagaatc	ctgaccaggt	gtacttggtg	gattatggcc	900
ttgcttatcg	gtactgcccc	gaaggagtgc	ataaagaata	caaagaagac	cccaaaagat	960
gtcacgatgg	cactattgaa	ttcacgagca	tcgatgcaca	caatgggtgtg	gccccatcaa	1020
gacgtggtga	tttggaataa	cttggttatt	gcgatgatcca	atggcttact	ggccatcttc	1080
cttggggagg	taatttgaaa	gattcctaat	atggttagaga	ttccaaaatt	agatacagag	1140
aaaatatttc	aagtttgatg	gacaaatggt	ttcctgagaa	aaacaaacca	ggtgaaaattg	1200
ccaaatacat	ggaaacagtg	aaattactag	actacactga	aaaacctctt	tatgaaaatt	1260
tacgtgacat	tcttttgcaa	ggactaaaag	ctataggaag	taaggatgat	ggcaaatggg	1320
acctcagtg	tgtggagaat	ggaggtttga	aagcaaaaac	aataacaaag	aagcgaaaga	1380
aagaaattga	agaaagcaag	gaacctgggtg	ttgaagatac	ggaatgggtca	aacacacaga	1440
cagaggaggc	catacagacc	cgttcaagaa	ccagaaaagag	agtccagaag	taattcagat	1500
gctgtgaacc	agatttcctt	ttctttgttt	tcttttgact	tttttctcct	tttctatttg	1560
aaactgtttt	ttttctgtg	agtcttgoga	ggtggaaagta	atgattaaat	actcatgtgt	1620
tcagaaaaca	taaacttttt	ttataaaaa	attttgtaca	attcattaaa	ggctaattta	1680
tgaaatttga	aaatcttcag	gttatactcc	ttaagttatc	ccaaagccgt	gtgtttgtga	1740

tggttttgag tacatatata tgaaaattat tatgacacgc acittttctaa tcattgtaca 1800  
 tttctcagag tggataaaaa tgtttgacaa agtcctcact ttttaaggaaa tgcaaagctt 1860  
 aaaataaaac tctcttttgc ttgatgc 1887

<210> 3  
 <211> 300  
 <212> PRT  
 <213> Homo sapiens

<220> -  
 <223> g222701

<400> 3  
 Met Asn Phe Gln Gly Leu Val Leu Thr Asp Asn Cys Lys Asn Gln  
 1 5 10 15  
 Trp Val Val Gly Pro Leu Ile Gly Lys Gly Gly Phe Gly Ser Ile  
 20 25 30  
 Tyr Thr Thr Asn Asp Asn Asn Tyr Val Val Lys Ile Glu Pro Lys  
 35 40 45  
 Ala Asn Gly Ser Leu Phe Thr Glu Gln Ala Phe Tyr Thr Arg Val  
 50 55 60  
 Leu Lys Pro Ser Val Ile Glu Glu Trp Lys Lys Ser His Asn Ile  
 65 70 75  
 Lys His Val Gly Leu Ile Thr Cys Lys Ala Phe Gly Leu Tyr Lys  
 80 85 90  
 Ser Ile Asn Val Glu Tyr Arg Phe Leu Val Ile Asn Arg Leu Gly  
 95 100 105  
 Ala Asp Leu Asp Ala Val Ile Arg Ala Asn Asn Asn Arg Leu Pro  
 110 115 120  
 Lys Arg Ser Val Met Leu Ile Gly Ile Glu Ile Leu Asn Thr Ile  
 125 130 135  
 Gln Phe Met His Glu Gln Gly Tyr Ser His Gly Asp Ile Lys Ala  
 140 145 150  
 Ser Asn Ile Val Leu Asp Gln Ile Asp Lys Asn Lys Leu Tyr Leu  
 155 160 165  
 Val Asp Tyr Gly Leu Val Ser Lys Phe Met Ser Asn Gly Glu His  
 170 175 180  
 Val Pro Phe Ile Arg Asn Pro Asn Lys Met Asp Asn Gly Thr Leu  
 185 190 195  
 Glu Phe Thr Pro Ile Asp Ser His Lys Gly Tyr Val Val Ser Arg  
 200 205 210  
 Arg Gly Asp Leu Glu Thr Leu Gly Tyr Cys Met Ile Arg Trp Leu  
 215 220 225  
 Gly Gly Ile Leu Pro Trp Thr Lys Ile Ser Glu Thr Lys Asn Cys  
 230 235 240  
 Ala Leu Val Ser Ala Thr Lys Gln Lys Tyr Val Asn Asn Thr Ala  
 245 250 255  
 Thr Leu Leu Met Thr Ser Leu Gln Tyr Ala Pro Arg Glu Leu Leu  
 260 265 270  
 Gln Tyr Ile Thr Met Val Asn Ser Leu Thr Tyr Phe Glu Glu Pro  
 275 280 285  
 Asn Tyr Asp Glu Phe Arg His Ile Leu Met Gln Gly Val Tyr Tyr  
 290 295 300







## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> C12N 9/12, 15/54, C07K 16/18, A61K 38/45, C12P 21/00, C12Q 1/68, C07K 16/40		<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 99/11794</b>
			<b>(43) International Publication Date:</b> 11 March 1999 (11.03.99)
<b>(21) International Application Number:</b> PCT/US98/18524		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
<b>(22) International Filing Date:</b> 4 September 1998 (04.09.98)		<b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
<b>(30) Priority Data:</b> 08/923,469 4 September 1997 (04.09.97) US		<b>(88) Date of publication of the international search report:</b> 8 July 1999 (08.07.99)	
<b>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application</b> US 08/923,469 (CIP) Filed on 4 September 1997 (04.09.97)			
<b>(71) Applicant (for all designated States except US):</b> INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US).			
<b>(72) Inventors; and</b>			
<b>(75) Inventors/Applicants (for US only):</b> YUE, Henry [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). LAL, Preeti [IN/US]; 2382 Lass Drive, Santa Clara, CA 95054 (US). CORLEY, Neil, C. [US/US]; 1240 Dale Avenue #30, Mountain View, CA 94040 (US).			
<b>(74) Agents:</b> BILLINGS, Lucy, J. et al.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US).			
<b>(54) Title:</b> NEW VRK1 KINASE			
<b>(57) Abstract</b> <p>The invention provides a new VRK1 kinase (NVRK1) and polynucleotides which identify and encode NVRK1. The invention also provides expression vectors, host cells, agonists, antibodies and antagonists. The invention also provides methods for treating disorders associated with expression of NVRK1.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## INTERNATIONAL SEARCH REPORT

In. ational Application No

PCT/US 98/18524

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N9/12 C12N15/54 C07K16/18 A61K38/45 C12P21/00  
C12Q1/68 C07K16/40

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	NEZU J. ET AL.: "VRK1, COMPLETE CDS" EMBL DATABASE, 1 May 1997, XP002096179 HEIDELBERG, DE Accession Number Q99986 ---	1-3, 5-11, 13, 18, 19
X, P	NEZU J. ET AL.: "Identification of two novel human putative serine/threonine kinases VRK1 and VR2 with structural similarity to vaccinia virus B1R kinase" GENOMICS, vol. 45, no. 2, 15 October 1997, pages 327-331, XP002096180 see the whole document --- -/--	1-12, 18, 19



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

27 April 1999

Date of mailing of the international search report

18/05/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Panzica, G

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/18524

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P	<p>WO 98 29552 A (CHUGAI RES INST MOLECULAR  MED ;NEZU JUN ICHI (JP); OKU ASUKA (JP))  9 July 1998  see abstract  see page 21 - page 24  see claim 1</p> <p style="text-align: center;">-----</p>	<p>1-12, 18,  19</p>

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/18524

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☒ Claims Nos.: 14-17  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
Remark: Claims 14-17, relating to agonists and antagonists for the purified protein kinase VRK1 of claim 1, could not be searched as their subject matter was not sufficiently disclosed.
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
  
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
  
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/18524

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9829552 A	09-07-1998	AU 5340698 A	31-07-1998